

Self-energy effects in the Polchinski and Wick-ordered renormalization-group approaches

A. Katanin

Institute of Metal Physics, 620041, Ekaterinburg, Russia

Abstract. I discuss functional renormalization group (fRG) schemes, which allow for non-perturbative treatment of the self-energy effects and do not rely on the one-particle irreducible functional. In particular, I consider Polchinskii or Wick-ordered schemes with amputation of full (instead of bare) Green functions, as well as more general schemes, and establish their relation to the ‘dynamical adjustment propagator’ scheme by M. Salmhofer [Ann. der Phys. 16, 171 (2007)]. While in the Polchinski scheme the amputation of full (instead of bare) Green functions improves treatment of the self-energy effects, the structure of the corresponding equations is not suitable to treat strong-coupling problems; it is not also evident, how the mean-field (MF) solution of these problems is recovered in this scheme. For Wick ordered scheme, excluding fully or partly tadpole diagrams one can obtain forms of fRG hierarchy, which are suitable to treat strong-coupling problems. In particular, I emphasize usefulness of the schemes, which are local in cutoff parameter, and compare them to the one-particle irreducible approach.

1. Introduction

Applications of functional renormalization group (fRG) approach [1, 2, 3] to problems of condensed matter and high energy physics have received substantial progress recently[6]. However, correct treatment of the self-energy effects remains one of the most difficult points of many fRG schemes.

Currently, the one-particle irreducible (1PI) scheme[3, 4, 5, 6, 7, 8, 9] is mostly used for treatment of the self-energy effects within fRG. Simple truncations within this scheme, however, do not allow to fulfill Ward identities and sometimes do not reproduce even the results of the mean-field approach, for which correct treatment of the self-energy effects is crucial. The truncation of 1PI equations of Ref. [10] allowed to reproduce the mean-field results and improve the results of more sophisticated approaches to fulfill Ward identities better. This approach found its applications in the treatment of weakly- and moderately interacting single-impurity problems[11], Hubbard model in the symmetry broken phase[12], and, more recently, two- and three dimensional Heisenberg model[13].

At the same time, 1PI approach has severe problems in describing the strong coupling regime of many-body systems. The problem of application of this approach in this case is mainly related to the one-loop structure of 1PI hierarchy of fRG equations. Although higher-loop contributions can be obtained by reinserting of the one-loop contributions from higher- to lower-order vertices, this procedure is usually spoiled by truncations of the hierarchy and the projection schemes used in approximate solutions, which do not allow getting correct results for non-ladder diagrams at a certain loop order[14]. In particular, the truncation, which neglects 8-point vertices and accounts for the full frequency- and momentum dependence of the 4-point and 6-point vertices (which is already a very complicated problem), yields a correct result for two-loop diagrams only [14].

On the other hand, some non-1PI schemes, in particular Wick-ordered scheme[2], already in the original formulation explicitly include many-loop contributions and can, in principle, improve treatment of the strong-coupling regime. The Wick-ordered scheme in its original formulation is however not suited to treat self-energy effects in the non-perturbative way. The change from amputation of the bare to interacting Green functions, proposed in Ref. [2], is not sufficient for such a treatment, since the function, with respect to which the Wick ordering is performed, should be also adjusted appropriately. A way for non-perturbative treatment of the self-energy effects was considered in the ‘dynamical adjustment’ scheme of Ref. [15]. In the present paper we propose somewhat different approach for non-perturbative treatment of the self-energy effects and consider in detail both, local schemes, which do not remove fully tadpole terms from the RG hierarchy, as well as non-local schemes, removing fully tadpole terms. We compare the approaches of the paper to the dynamical adjusting scheme of Ref. [15] and the 1PI approach, as well as discuss the results of the application of these methods to the toy model.

2. Self-energy effects in the Polchinskii scheme

We consider the model described by an action

$$\mathcal{S}_\Lambda[\bar{\psi}, \psi] = \mathcal{S}_0^\Lambda[\bar{\psi}, \psi] + \mathcal{V}[\bar{\psi}, \psi], \quad (1)$$

where $\bar{\psi}, \psi$ are the bosonic or fermionic (Grassman) fields, $\mathcal{V}$ is an interaction and

$$\mathcal{S}_0^\Lambda[\bar{\psi}, \psi] = \int d^d x \int d\tau \bar{\psi} (C_{0,\Lambda})^{-1} \psi \quad (2)$$

is quadratic in fields. $C_{0,\Lambda}$ is the cut bare propagator, e.g. having the Fourier transform $C_{0,\Lambda} = \chi_{>,\Lambda} / (i\omega_n - \varepsilon_{\mathbf{k}})$, where $i\omega_n$ are bosonic or fermionic Matsubara frequencies, $\varepsilon_{\mathbf{k}}$ is the dispersion, and $\chi_{>,\Lambda}$ is the cutoff function, which cuts off the low-energy modes, e.g. $\chi_{>,\Lambda_0} = 0$ and $\chi_{>,0} = 1$.

The standard procedure of derivation of fRG equations relies on the differentiating of generating functional for connected vertices, amputated with the non-interacting Green functions,

$$\mathcal{V}_\Lambda[\bar{\psi}, \psi] = -\ln \int D[\bar{\chi}, \chi] e^{-\mathcal{S}_0^\Lambda[\bar{\chi}, \chi] - \mathcal{V}[\bar{\psi} + \bar{\chi}, \psi + \chi]} \quad (3)$$

This procedure can be supplemented by a consequent reamputation of external legs, changing their amputation from non-interacting to that with interacting Green functions.

For the purposes of the present paper, we consider somewhat more general procedure, which allows to obtain the renormalization-group equations accounting for the self-energy effects with further possible generalizations. To this end, we introduce the counterterm $\bar{\Sigma}_\Lambda \bar{\psi} \psi$ into the bare action and replace $(C_{0,\Lambda})^{-1} \rightarrow C_\Lambda^{-1}$:

$$\mathcal{S}_\Lambda[\bar{\psi}, \psi] = \bar{\psi} C_\Lambda^{-1} \psi + \bar{\psi} \bar{\Sigma}_\Lambda \psi + \mathcal{V}[\bar{\psi}, \psi] \quad (4)$$

Although for the choice

$$C_\Lambda^{-1} = (C_{0,\Lambda})^{-1} - \bar{\Sigma}_\Lambda \quad (5)$$

the counterterm is cancelled, we will not in general assume validity of equation (5) during the flow, requiring its fulfillment only in the end of the flow. We also treat $\bar{\Sigma}_\Lambda$ as a part of the interaction, such that amputation of the vertices by the functions C_Λ is achieved naturally in this way, since the non-interacting term now contains C_Λ^{-1} . Introducing the effective interaction in the standard way

$$e^{-\mathcal{V}_\Lambda[\bar{\eta}, \eta]} = \int D[\psi, \bar{\psi}] e^{-\bar{\psi} (C_\Lambda^{-1}) \psi - \mathcal{V}[\bar{\psi} + \bar{\eta}, \psi + \eta] + \bar{\Sigma}_\Lambda (\bar{\psi} + \bar{\eta})(\psi + \eta)} \quad (6)$$

and performing the same steps, as for deriving Polchinski equation, we obtain

$$\partial_\Lambda \mathcal{V}_\Lambda = \Delta_{\dot{C}_\Lambda} \mathcal{V}_\Lambda - \Delta_{\dot{C}_\Lambda}^{12} \mathcal{V}_\Lambda^{(1)} \mathcal{V}_\Lambda^{(2)} + \dot{\bar{\Sigma}}_\Lambda \frac{\delta \mathcal{V}_\Lambda}{\delta \bar{\Sigma}_\Lambda} \quad (7)$$

where $\Delta_{\dot{C}_\Lambda} \mathcal{V}_\Lambda = \text{Tr}(\dot{C}_\Lambda \delta^2 \mathcal{V}_\Lambda / \delta^2 \Psi)$, $\Delta_{\dot{C}_\Lambda}^{12} \mathcal{V}_\Lambda^{(1)} \mathcal{V}_\Lambda^{(2)} = \left(\frac{\delta \mathcal{V}_\Lambda}{\delta \Psi}, \dot{C}_\Lambda \frac{\delta \mathcal{V}_\Lambda}{\delta \Psi} \right)$, $\Psi = (\psi, \bar{\psi})$, ∂_Λ and dot stand for the derivative over Λ . Expressing the variational derivative $(\delta \mathcal{V}_\Lambda / \delta \bar{\Sigma}_\Lambda)$

through variational derivatives of $\mathcal{V}_\Lambda$ over fields, we obtain

$$\begin{aligned} \partial_\Lambda(\mathcal{V}_\Lambda - \bar{\Sigma}_\Lambda \bar{\psi}\psi) &= \Delta_{\dot{C}_\Lambda - \dot{\bar{\Sigma}}_\Lambda C^2} \mathcal{V}_\Lambda - \Delta_{\dot{C}_\Lambda - \dot{\bar{\Sigma}}_\Lambda C^2 \Lambda}^{12} \mathcal{V}_\Lambda^{(1)} \mathcal{V}_\Lambda^{(2)} \\ &\quad - \dot{\bar{\Sigma}}_\Lambda C \left(\bar{\psi} \frac{\delta \mathcal{V}_\Lambda}{\delta \bar{\psi}} + \psi \frac{\delta \mathcal{V}_\Lambda}{\delta \psi} \right) \end{aligned} \quad (8)$$

Choosing $\bar{\Sigma}_\Lambda = 0$, we arrive at the hierarchy of fRG equations in the Polchinski scheme[1]

$$\partial_\Lambda \mathcal{V}_\Lambda = \Delta_{\dot{C}_{0,\Lambda}} \mathcal{V}_\Lambda - \Delta_{\dot{C}_{0,\Lambda}}^{12} \mathcal{V}_\Lambda^{(1)} \mathcal{V}_\Lambda^{(2)}$$

Considering expansion of $\mathcal{V}_\Lambda$ in fields

$$\mathcal{V}_\Lambda = \sum_m V_m \Psi(x_1) \dots \Psi(x_m), \quad (9)$$

this hierarchy reads

$$\begin{aligned} \dot{V}_2 &= -V_2 \dot{C}_{0,\Lambda} V_2 + V_4 \circ \dot{C}_{0,\Lambda}, \\ \dot{V}_4 &= -4V_2 \dot{C}_{0,\Lambda} V_4 + V_6 \circ \dot{C}_{0,\Lambda}, \\ \dot{V}_6 &= -6V_2 \dot{C}_{0,\Lambda} V_6 + V_4 \dot{C}_{0,\Lambda} V_4 + V_8 \circ \dot{C}_{0,\Lambda}, \\ &\dots, \end{aligned} \quad (10)$$

$\circ$ denotes convolution with respect to internal momenta and frequencies. On the other hand, for the choice $V_2 = 0$ (8) yields equations for the self-energy $\Sigma_\Lambda = \bar{\Sigma}_\Lambda$ and fully amputated vertex functions, which are denoted in the following as H_m :

$$\begin{aligned} \dot{\Sigma}_\Lambda &= -H_4 \circ (\dot{C}_\Lambda - \dot{\Sigma}_\Lambda C^2), \\ \dot{H}_4 &= H_6 \circ (\dot{C}_\Lambda - \dot{\Sigma}_\Lambda C^2) - 4H_4 C_\Lambda \dot{\Sigma}_\Lambda, \\ \dot{H}_6 &= H_8 \circ (\dot{C}_\Lambda - \dot{\Sigma}_\Lambda C^2) + H_4 (\dot{C}_\Lambda - \dot{\Sigma}_\Lambda C^2) H_4 - 6H_6 C_\Lambda \dot{\Sigma}_\Lambda, \\ &\dots \end{aligned} \quad (11)$$

Assuming that the relation (5) between the full and bare Green functions is fulfilled during the flow, the scheme (11) can be obtained from equations (10) by substituting

$$\begin{aligned} V_2 &= H_2 (C_\Lambda / C_{0,\Lambda})^2 = - (C_\Lambda / C_{0,\Lambda}) \Sigma_\Lambda \\ V_4 &= H_4 (C_\Lambda / C_{0,\Lambda})^4, \quad \dots \end{aligned} \quad (12)$$

For the choice (5) we also have $\dot{C}_\Lambda - \dot{\Sigma}_\Lambda C_\Lambda^2 = \dot{C}_{0,\Lambda} / (1 - C_{0,\Lambda} \Sigma_\Lambda)^2$, which allows to recover the equations for amputated vertices, derived in Ref. [2]. However, as it was already mentioned, general equations (11) do not assume the validity of the relation (5) during the flow; they are valid for any functional $C_\Lambda[\Sigma_\Lambda]$, which fulfills $C_{\Lambda=\Lambda_0} = 0$ in the beginning of the flow and the relation (5) in the end of the flow.

The equation for the self-energy (11) is similar to its form in 1PI scheme (and coincides with this scheme for the choice (5)). However, the equations for higher-order vertices are organized differently: the pairs of vertices are connected only by tree-like diagrams, while one- and higher loop contributions are obtained by substituting results from higher- to lower order vertices, which may lead to difficulties in the strong-coupling regime. These equations also do not allow to see easily how the mean-field results are reproduced.

3. Wick-ordered schemes

3.1. General consideration and non-local schemes

To overcome these difficulties, we following to Refs. [2, 15] consider the Wick ordered modification of Eqs. (10). This scheme considers an expansion of the effective action in the Wick-ordered monomials $\Omega_{D_\Lambda}(\Psi(x_1) \dots \Psi(x_m))$

$$\mathcal{V}_\Lambda = \sum_m W_m \Omega_{D_\Lambda}(\Psi(x_1) \dots \Psi(x_m)) = e^{-\Delta_{D_\Lambda}} \tilde{\mathcal{V}}_\Lambda \quad (13)$$

$$\tilde{\mathcal{V}}_\Lambda = \sum_m W_m \Psi(x_1) \dots \Psi(x_m) \quad (14)$$

where the Wick-ordering propagator D_Λ fulfills $D_{\Lambda=0} = 0$. Performing Wick-ordering of equations (8) using (13) and assuming $W_2 = 0$, we obtain

$$\begin{aligned} \dot{\Sigma}_\Lambda &= - \left[\frac{d(D_\Lambda + C_\Lambda)}{d\Lambda} - \dot{\Sigma}_\Lambda C_\Lambda (2D_\Lambda + C_\Lambda) \right] H_4 \\ &\quad + \sum_{m_1, m_2 \geq 4} H_{m_1} \circ (\dot{C}_\Lambda - \dot{\Sigma}_\Lambda C_\Lambda^2) \circ D_\Lambda^{\frac{m_1+m_2}{2}-2} \circ H_{m_2} \\ \dot{H}_m &= \left[\frac{d(D_\Lambda + C_\Lambda)}{d\Lambda} - \dot{\Sigma}_\Lambda C_\Lambda (2D_\Lambda + C_\Lambda) \right] H_{m+2} - m H_m \dot{\Sigma}_\Lambda C_\Lambda \\ &\quad - \sum_{m_1, m_2 \geq 4} H_{m_1} \circ (\dot{C}_\Lambda - \dot{\Sigma}_\Lambda C_\Lambda^2) \circ D_\Lambda^{\frac{m_1+m_2-m}{2}-1} \circ H_{m_2} \quad (m \geq 4) \end{aligned} \quad (15)$$

In the square brackets we have grouped together terms, corresponding to generalised tadpole diagrams. Standard choice [2] is $\dot{D}_\Lambda = -\dot{C}_{\Lambda,0}$. Let however choose D such that tadpole diagrams, including the contributions proportional to $\dot{\Sigma}$, cancel out. Then we have the following differential equation for D :

$$\frac{d(D_\Lambda + C_\Lambda)}{d\Lambda} - \dot{\Sigma}_\Lambda C_\Lambda (2D_\Lambda + C_\Lambda) = 0 \quad (16)$$

The resulting hierarchy of equations is simplified and read

$$\begin{aligned} \dot{\Sigma}_\Lambda &= \sum_{m_1, m_2 \geq 4} H_{m_1} \circ (\dot{C}_\Lambda - \dot{\Sigma}_\Lambda C_\Lambda^2) \circ D_\Lambda^{\frac{m_1+m_2}{2}-2} \circ H_{m_2} \\ \dot{H}_m &= -m H_m \dot{\Sigma}_\Lambda C_\Lambda \\ &\quad - \sum_{m_1, m_2 \geq 4} H_{m_1} \circ (\dot{C}_\Lambda - \dot{\Sigma}_\Lambda C_\Lambda^2) \circ D_\Lambda^{\frac{m_1+m_2-m}{2}-1} \circ H_{m_2} \quad (m \geq 4) \end{aligned} \quad (17)$$

Similarly to Ref. [15], to guarantee that $D_{\Lambda=0} = 0$ one has to solve Eq. (16) in terms of corresponding condition at $\Lambda = 0$, instead of using initial condition at Λ_0 . The corresponding solution to the equation (16) has the form

$$D_\Lambda = - \int_0^\Lambda d\Lambda' (\dot{C}_{\Lambda'} - \dot{\Sigma}_{\Lambda'} C_{\Lambda'}^2) \exp \left(2 \int_{\Lambda'}^\Lambda d\Lambda'' C_{\Lambda''} \dot{\Sigma}_{\Lambda''} \right) \quad (18)$$

In particular, for the choice of the propagator (5) we obtain

$$D_\Lambda = \int_0^\Lambda d\Lambda' \dot{D}_{\Lambda',0} \exp \left(2 \int_{\Lambda'}^\Lambda d\Lambda'' C_{\Lambda''} \dot{\Sigma}_{\Lambda''} \right)$$

$$= \frac{1}{(1 - C_{0,\Lambda}\Sigma_\Lambda)^2} \int_0^\Lambda d\Lambda' \dot{C}_{0,\Lambda}' \exp \left(2 \int_{\Lambda'}^\Lambda d\Lambda'' \frac{\Sigma_{\Lambda''} \dot{C}_{0,\Lambda}'}{1 - C_{0,\Lambda''}\Sigma_{\Lambda''}} \right) \quad (19)$$

and

$$\begin{aligned} \dot{\Sigma}_\Lambda &= \frac{1}{2} \sum_{m_1, m_2 \geq 4} H_{m_1} \circ \frac{1}{(1 - C_{0,\Lambda}\Sigma_\Lambda)^2} \dot{C}_{0,\Lambda}' \circ D_\Lambda^{\frac{m_1+m_2}{2}-2} \circ H_{m_2} \\ \dot{H}_m &= -m H_m C_\Lambda \dot{\Sigma}_\Lambda \\ &\quad - \frac{1}{2} \sum_{m_1, m_2 \geq 4} H_{m_1} \circ \frac{1}{(1 - C_{0,\Lambda}\Sigma_\Lambda)^2} \dot{C}_{0,\Lambda}' \circ D_\Lambda^{\frac{m_1+m_2-m}{2}-1} \circ H_{m_2} \end{aligned} \quad (20)$$

where $\dot{C}_{0,\Lambda}' = -\dot{C}_{0,\Lambda}$ and $\dot{D}_{\Lambda,0} = \dot{C}_{0,\Lambda}'/(1 - C_\Lambda^0 \Sigma_\Lambda)^2$.

Let us see, how the mean-field solution is recovered from the equations (15) or (20). Since there are no tadpole diagrams for $d\Sigma_\Lambda/d\Lambda$, we have $d\Sigma_\Lambda^{\text{MF}}/d\Lambda = 0$, i.e. $\Sigma = \Sigma^{\text{MF}}$ is constant in Λ and equal to its mean-field value, which is given by a (self-consistently determined) sum of tadpole diagrams, absorbed into the definition of Wick ordering. Equation (18) is easily integrated and yields

$$D_\Lambda^{\text{MF}} = C_{\Lambda=0}^{\text{MF}} - C_\Lambda^{\text{MF}} \quad (21)$$

In case of fermions and moving Fermi surface ($\text{Re}\Sigma \neq 0$) this function for smooth cutoff and intermediate Λ has singularity at both, physical and running Fermi surface. Since $\dot{D}_\Lambda^{\text{MF}} = -\dot{C}_\Lambda^{\text{MF}}$, the equation for H_4 can be then also solved analytically to reproduce the standard random-phase approximation (RPA) result

$$H_4 = \frac{H_4^{\text{MF}}}{1 - H_4^{\text{MF}} \text{Tr}(C_\Lambda^{\text{MF}})^2}. \quad (22)$$

Returning to the treatment of the interaction beyond mean-field theory, the advantage of the equations (19) and (20) is in their simple form in the sharp cutoff limit. Assuming $\chi_{>,\Lambda} = \theta(|\varepsilon_{\mathbf{k}}| - \Lambda)$, $C_{0,\Lambda} = \chi_{>,\Lambda}/(i\omega_n - \varepsilon_{\mathbf{k}})$, we obtain

$$D_\Lambda = \frac{\theta(\Lambda - |\varepsilon_{\mathbf{k}}|)}{i\omega_n - \varepsilon_{\mathbf{k}} - \Sigma_{\Lambda'=|\varepsilon_{\mathbf{k}}|}} \quad (23)$$

which requires the knowledge of $\Sigma_{\Lambda'}$ at $\Lambda' < \Lambda$, which has to be determined self-consistently and the single-scale propagator

$$F_\Lambda = \frac{\delta(\Lambda - |\varepsilon_{\mathbf{k}}|)}{i\omega_n - \varepsilon_{\mathbf{k}} - \Sigma_\Lambda}, \quad (24)$$

Both propagators have simpler form, than those in Ref. [15], where additional integration over the cutoff parameter is still present even for the sharp cutoff (see Appendix).

The equations (15) can be somewhat generalized by considering reamputation of vertices $H_\Lambda^{(n)} = \widetilde{H}_\Lambda^{(n)}(S_\Lambda)^n$ with respect to the amputation by full Green functions C_Λ (corresponding to $S_\Lambda = 1$),

$$S_\Lambda^{-2} \dot{\Sigma}_\Lambda = - \left[\frac{d\widetilde{D}_\Lambda}{d\Lambda} - 2\widetilde{D}_\Lambda (\dot{\Sigma}_\Lambda C_\Lambda + \dot{S}_\Lambda S_\Lambda^{-1}) + S_\Lambda^2 (\dot{C}_\Lambda - \dot{\Sigma}_\Lambda C_\Lambda^2) \right] \circ \widetilde{H}_4$$

$$\begin{aligned}
 & + \sum_{m_1, m_2 \geq 4} \widetilde{H}_{m_1} \circ S_\Lambda^2 (\dot{C}_\Lambda - \dot{\Sigma}_\Lambda C_\Lambda^2) \circ \widetilde{D}_\Lambda^{\frac{m_1+m_2}{2}-2} \circ \widetilde{H}_{m_2} \\
 \dot{\widetilde{H}}_m & = \left[\frac{d\widetilde{D}_\Lambda}{d\Lambda} - 2\widetilde{D}_\Lambda (\dot{\Sigma}_\Lambda C_\Lambda + \dot{S}_\Lambda S_\Lambda^{-1}) + S_\Lambda^2 (\dot{C}_\Lambda - \dot{\Sigma}_\Lambda C_\Lambda^2) \right] \circ \widetilde{H}_{m+2} \\
 & - \sum_{m_1, m_2 \geq 4} \widetilde{H}_{m_1} \circ S_\Lambda^2 (\dot{C}_\Lambda - \dot{\Sigma}_\Lambda C_\Lambda^2) \circ \widetilde{D}_\Lambda^{\frac{m_1+m_2-m}{2}-1} \circ \widetilde{H}_{m_2} \\
 & - m \widetilde{H}_m (C_\Lambda \dot{\Sigma}_\Lambda + \dot{S}_\Lambda S_\Lambda^{-1})
 \end{aligned} \tag{25}$$

where $\widetilde{D}_\Lambda = D_\Lambda S_\Lambda^2$. Equations (25) allow to establish connection with the dynamic adjustment propagator scheme of Ref. [15]. The explicit relation between the equations (25) and those of Ref. [15] is discussed in the Appendix and involves passing from the function C_Λ to a different function D_Λ^M with the use of equation (37).

If, similarly to Ref. [15], we require $C_\Lambda \dot{\Sigma}_\Lambda + \dot{S}_\Lambda S_\Lambda^{-1} = 0$, then equations (25) take simpler form due to cancellation of some terms in this specially chosen amputation of vertices. This way, equations (25) allow also to establish simpler view on the result (18). Indeed, let assume that $\dot{\Sigma}_\Lambda C_\Lambda + \dot{S}_\Lambda S_\Lambda^{-1} = 0$ for some S . Then we have

$$S_\Lambda = \exp \left(\int_\Lambda^{\Lambda_0} \dot{\Sigma}_{\Lambda'} C_{\Lambda'} d\Lambda' \right) \tag{26}$$

The condition of vanishing generalized tadpole diagrams yields $\frac{d\widetilde{D}_\Lambda}{d\Lambda} = -S_\Lambda^2 (\dot{C}_\Lambda - \dot{\Sigma}_\Lambda C_\Lambda^2)$, which leads us to equation (18). Although the condition (18) generally does not yield Wick propagators, which derivative coincides with the single-scale propagator $\dot{\Sigma}_\Lambda C_\Lambda^2 - \dot{C}_\Lambda$, this is achieved in the amputated scheme with the amputation factor S_Λ given by the equation (26). In general, analytical or numerical evaluation of the integral in this equation can be, however, rather complicated.

3.2. Local schemes

In practical applications, the schemes, which are local in Λ , may have some advantage. Simplest choice of the propagators

$$D_\Lambda = \frac{\chi_{<,\Lambda}}{C_0^{-1} - \Sigma_\Lambda}; C_\Lambda = \frac{\chi_{>,\Lambda}}{C_0^{-1} - \Sigma_\Lambda} \tag{27}$$

suffers from the problem, discussed in Ref. [15], namely it yields remaining tadpole terms in $\partial_\Lambda H_m$, proportional to $(D_\Lambda^2 \partial_\Lambda \Sigma_\Lambda) \circ H_{m+2}$, which have potential infrared divergencies due to square of the propagator D_Λ . To avoid this problem, we choose the propagator C_Λ according to the equation (5) and use similar expression for D_Λ , including the mean-field self-energy into the bare propagator and choosing non-interacting part of the action $C_{0,\Lambda} = \chi_{>,\Lambda} \overline{C}_{0,\Lambda}$,

$$D_\Lambda = \frac{\chi_{<,\Lambda}}{\overline{C}_{0,\Lambda}^{-1} - \Sigma^{\text{MF}} - \chi_{>,\Lambda} \Sigma_\Lambda}; C_\Lambda = \frac{\chi_{>,\Lambda}}{\overline{C}_{0,\Lambda}^{-1} - \Sigma^{\text{MF}} - \chi_{>,\Lambda} \Sigma_\Lambda}, \tag{28}$$

to obtain from the equations (15) the local flow equations

$$\dot{\Sigma}_\Lambda = -T_\Lambda \circ H_4 \tag{29}$$

$$\begin{aligned}
 & - \sum_{m_1, m_2 \geq 4} H_{m_1} \circ F_\Lambda \circ D_\Lambda^{\frac{m_1+m_2}{2}-2} \circ H_{m_2}, \\
 \dot{H}_m &= T_\Lambda \circ H_{m+2} - m H_m C_\Lambda \dot{\Sigma}_\Lambda \\
 & + \sum_{m_1, m_2 \geq 4} H_{m_1} \circ F_\Lambda \circ D_\Lambda^{\frac{m_1+m_2-m}{2}-1} \circ H_{m_2}.
 \end{aligned}$$

where

$$\begin{aligned}
 T_\Lambda &= \frac{r \bar{C}_{0,\Lambda}^{-2} \dot{\bar{C}}_{0,\Lambda} \chi_{>,\Lambda} + r \Sigma_\Lambda \dot{\chi}_{>,\Lambda} - \chi_{<,\Lambda} \chi_{>,\Lambda} \dot{\Sigma}_\Lambda}{(\bar{C}_{0,\Lambda}^{-1} - \Sigma^{\text{MF}} - \chi_{>,\Lambda} \Sigma_\Lambda)^2} \\
 F_\Lambda &= - \frac{\bar{C}_{0,\Lambda}^{-2} \dot{\bar{C}}_{0,\Lambda} \chi_{>,\Lambda} + (\bar{C}_{0,\Lambda}^{-1} - \Sigma^{\text{MF}}) \dot{\chi}_{>,\Lambda}}{(\bar{C}_{0,\Lambda}^{-1} - \Sigma^{\text{MF}} - \chi_{>,\Lambda} \Sigma_\Lambda)^2}
 \end{aligned} \tag{30}$$

and we have assumed $\chi_{>,\Lambda} + \chi_{<,\Lambda} = r$. Due to special choice of propagators, the solution to the equations (29) has natural physical interpretation even at the intermediate stages of the flow, since it corresponds to the flow of the functional with the bare propagator $\chi_{>,\Lambda}/(C_{0,\Lambda}^{-1} - \Sigma^{\text{MF}})$ after the Wick ordering with the propagator D_Λ . The equations (29) have to be solved with the initial condition $\Sigma_\Lambda = 0$, since the constant (mean-field) initial part of Σ_Λ enters propagators explicitly through Σ^{MF} .

The scheme (28)-(30) may be useful for the realization of the interaction [8] and temperature- [5, 7, 9] flow in the Wick-ordered scheme. Indeed, the choice $\chi_{>,\Lambda} = \Lambda^{1/2}$, $\chi_{<,\Lambda} = 1 - \Lambda^{1/2}$, and $\bar{C}_{0,\Lambda} = C_0$ in the equation (27) is analogous to the interaction flow [8], while the choice $\Lambda = T$, $\chi_{>,T} = T^{-1/2}$, $r = T_0^{-1/2}$, and $\bar{C}_{0,T}^{-1} = (i\omega_n - \varepsilon_{\mathbf{k}})/T$, where T is the temperature and T_0 is the final temperature of the flow, is analogous to the temperature-flow in the 1PI approach [5, 7, 9]. The disadvantage of the latter scheme in Wick-ordered approach is that similarly to the 1PI flow with counterterms [16] it requires the final temperature of the flow as an input, and therefore does not allow to obtain the whole temperature evolution of the system for the same input parameters.

Within similar truncations of the hierarchy, the momentum-, frequency, or interaction flow in the Wick-ordered scheme with the choice of propagators (28) can be already superior to the 1PI scheme, since the latter contains tadpole terms of the order $O(H_{m+2})$ in the weak-coupling regime, which are needed to generate multiple-loop contributions from fRG hierarchy. On contrary, the Wick-ordered scheme contains these contributions explicitly, while the “dangerous” tadpole contributions have smaller value; the initial (mean-field) self-energy is also already included in both propagators C_Λ and D_Λ . Specifically, in the weak-coupling limit in the scheme (29) we obtain $\Sigma_\Lambda \sim \dot{\Sigma}_\Lambda = O(H_4^2)$, yielding therefore the correction to $\dot{\Sigma}_\Lambda$ from the remaining tadpole diagrams of the order $O(H_4^3)$ and the corresponding correction to the vertices $H_{m \geq 4}$ of the order $O(H_4^2 H_{m+2})$.

Smaller value of the tadpole terms in Wick-ordered scheme and explicit presence of multi-loop contributions also implies possibility of having less fine parametrization of vertices, required for the solution of the equations (29) and providing the same accuracy, as in 1PI scheme. Indeed, the effort of extra evaluation of each loop in Wick ordered

scheme scales as $n_p n_o$ where n_p is the number of integration points in momentum-frequency space and n_o is the number of internal (e.g. spin, orbital, etc.) degrees of freedom. At the same time, finer parametrization of m -point vertex in 1PI scheme increases computational effort by $(n_1/n_2)^{m-1}$ where n_1 and $n_2 < n_1$ are the number of patches in momentum-frequency space for the 1PI and Wick-ordered equations, respectively. For $n_1 \sim n_p/10$, $n_2 \sim 10$ and for not very large number of internal degrees of freedom, the increase of the effort in 1PI scheme $(n_1/n_2)^{m-1}$, required to reproduce accurately multi-loop contributions, can easily exceed $n_p n_o$ even for the $m = 4$ -point vertex due to extra factor $n_p^2/(10^6 n_o)$, which is rather large in the practical applications, because of large $n_p \sim 10^4$.

Application of the full cancellation of the tadpole diagrams, given by the equation (19) or dynamical adjusting scheme of Ref. [15] may further improve applicability of the Wick-ordered equations, but is considerably more complicated because of the necessity to fulfill the self-consistency condition $D_{\Lambda=0}^{\text{phys}} = 0$ (or $R_{\Lambda=0} = 0$), which results in the requirement to know the self-energy $\Sigma_{\Lambda'}$ at the stages of the flow, which are later than the current one. Yet, even in this case one can search for easier integrable functions by adjusting C_{Λ} in equations (15), (25).

3.3. Toy model

To gain insight into the applicability of the described approaches, we consider the toy φ^4 -like model[17]

$$S_0 = \frac{a}{2}\varphi^2; \quad \mathcal{V} = \frac{b}{4!}\varphi^4 \quad (31)$$

In the following we put $a = 1/\chi_{>,\Lambda}$ with $\chi_{>,\Lambda} = 1 - \Lambda$ and use Λ as a scaling parameter, which changes from one (bare model) to zero. In the truncation $H_6 = 0$ the flow equations (16), (20) take the form

$$\begin{aligned} \dot{\Sigma}_{\Lambda} &= -\frac{1}{2}b_{\Lambda}^2 F_{\Lambda} D_{\Lambda}^2 \\ \dot{b}_{\Lambda} &= 3b_{\Lambda}^2 F_{\Lambda} D_{\Lambda} - 4b_{\Lambda} C_{\Lambda} \dot{\Sigma}_{\Lambda} \\ \dot{D}_{\Lambda} &= F_{\Lambda} + 2\dot{\Sigma}_{\Lambda} C_{\Lambda} D_{\Lambda} \end{aligned} \quad (32)$$

where we choose, according to the equation (28),

$$C_{\Lambda} = \frac{1 - \Lambda}{1 - \Sigma^{\text{MF}} - (1 - \Lambda)\Sigma_{\Lambda}}, \quad (33)$$

such that

$$F_{\Lambda} = -\dot{C}_{\Lambda} + \dot{\Sigma}_{\Lambda} C_{\Lambda}^2 = \frac{1 - \Sigma^{\text{MF}}}{(1 - \Sigma^{\text{MF}} - (1 - \Lambda)\Sigma_{\Lambda})^2}; \quad (34)$$

the initial conditions $\Sigma^{\text{MF}} = \frac{1}{2}(1 - \sqrt{1 + 2b})$, $\Sigma_{\Lambda=1} = 0$, $b_{\Lambda=1} = b$ and $D_{\Lambda=0} = 0$. In numerical calculations, we adjust the initial value $D_{\Lambda=1}$ to achieve vanishing of the propagator in the end of the flow.

On the other hand, the local equations (28), (29) yield

$$\begin{aligned}\dot{\Sigma}_\Lambda &= -\frac{1}{2}b_\Lambda T_\Lambda - \frac{1}{2}b_\Lambda^2 F_\Lambda D_\Lambda^2 \\ \dot{b}_\Lambda &= -4b_\Lambda C_\Lambda \dot{\Sigma}_\Lambda + 3b_\Lambda^2 F_\Lambda D_\Lambda\end{aligned}\tag{35}$$

with

$$\begin{aligned}D_\Lambda &= \frac{\Lambda}{1 - \Sigma^{\text{MF}} - (1 - \Lambda)\Sigma_\Lambda} \\ T_\Lambda &= \frac{\Sigma_\Lambda - \Lambda(1 - \Lambda)\dot{\Sigma}_\Lambda}{(1 - \Sigma^{\text{MF}} - (1 - \Lambda)\Sigma_\Lambda)^2}\end{aligned}\tag{36}$$

The results of the solution of equations (32) and (35) for the self-energy and coupling constant in the end of the flow for interaction strength $b = 1, 2, 3$ are presented in the Table. For comparison, we also present the result of 1PI approach [17] and the local scheme, suggested in Ref. [15], see also the equations (43) of the Appendix; we find numerically, that using the choice (27) of propagators in the equations (35) yields the same results, as the latter scheme. Note that, as discussed in previous Section, the local scheme based on the choice (27), as well as the local scheme of Ref. [15], are actually not suitable for treatment of many-body systems because of possible divergencies in the flow equations. The non-local (adjusting) scheme of Ref. [15] (see also equations (41) of the Appendix) is found to yield numerically very close results to the equations (32)-(34).

One can see from the Table, that for the model (31) all considered Wick-ordered schemes yield improvement in comparison to 1PI scheme; the equations (35) improve the results for vertices of the local approach of Ref. [15], the self-energies of the two approaches are practically identical. Surprisingly, being computationally more expensive and yielding slight improvement of 1PI scheme, the approaches, fully excluding tadpole diagrams, yield worse results for the considering toy model, than the local schemes. Therefore, the adjustment of the Wick propagator to exclude fully tadpole diagrams does not generally implies smaller truncation errors. This fact can be attributed to the property of the Wick-ordered equations in general amputation scheme (25) to have derivative of the Wick propagator non-equal to the single-scale propagator. Therefore, some remaining tadpole terms seem necessary to compensate this difference. Note that choosing special amputation of the vertices, which provide equality of the derivative of the Wick propagator to the single-scale propagator, such as given by the Eq. (26) or considered in Ref. [15], does not yield further improvement of the truncation, since the amputation itself does not change physically observable quantities. For more complicated problems, one may also expect that the schemes, which are local in the cutoff parameter, but do not exclude fully tadpole diagrams (such as the scheme (29)) are more preferable.

Method	$-\Sigma$			b_R		
$b =$	1	2	3	1	2	3
1PI, Ref. [17]	0.320398	0.519824	0.677443	0.513382	0.809275	1.05686
Eqs. (17), Eqs. (32)	0.323881	0.527393	0.687805	0.530763	0.855922	1.13321
local, Ref.[15]	0.330197	0.546350	0.720883	0.535013	0.874050	1.17181
Eqs.(29);Eqs.(35)	0.329972	0.545415	0.719051	0.542277	0.899275	1.21913
Exact	0.332425	0.550557	0.726505	0.607899	1.055430	1.46469

Table. The self-energy and connected fully amputated 4-point vertex b_R of the model (31) obtained by different methods for values of the coupling constant $b = 1, 2, 3$.

4. Summary

In Summary, we have considered the renormalization-group schemes, which allow to treat self-energy effects in a non-perturbative way and do not rely on using one-particle irreducible functionals. In the Wick-ordered scheme the choice of the propagator, with respect to which the Wick-ordering is performed, allowing to remove partly or fully tadpole diagrams from renormalization-group equations, has some advantage over 1PI formalism. While full removing of tadpole terms yields the differential equation with “finite” instead of initial condition, and the results, which are similar to the earlier proposed approach [15], removing partly tadpole terms we have obtained the differential equations, which are local with respect to the cutoff parameter and avoid potential divergencies in the flow equations. We have also argued that in general the suggested local approach may be superior to 1PI formalism, due to smaller contribution of the tadpole diagrams, which implies that in general it requires less fine parametrization of the vertices, than the 1PI approach. Further investigations of the necessity to keep some tadpole contributions in the Wick-ordering hierarchy and comparison of 1PI and Wick-ordered schemes for more complicated models would be of certain interest.

Acknowledgements. The author is grateful to M. Salmhofer for discussions on the properties of Wick-ordered approach and hospitality during the visit to the Institute of Theoretical Physics (Heidelberg), and also to Max-Planck Society for partial financial support within the Partnership Program.

Appendix. Relation to the dynamic adjustment scheme of Ref. [15].

In this Appendix we consider relation of the approach of present paper to that of Ref. [15]. To relate the equations (15) to those of Ref. [15], we put

$$C_\Lambda S^2 = \mathcal{C}_\Lambda - D_\Lambda^M \quad (37)$$

with some function D_Λ^M , which satisfies $D_{\Lambda=\Lambda_0}^M = \mathcal{C}_{\Lambda=\Lambda_0}$. Choosing $\mathcal{C} = S^2/(C_0^{-1} - \Sigma_\Lambda)$, we obtain

$$S^2(\dot{C}_\Lambda - \dot{\Sigma}_\Lambda C_\Lambda^2) = S^2\left(\frac{d}{d\Lambda}(\mathcal{C}/S^2) - \dot{\Sigma}_\Lambda \mathcal{C}^2/S^4\right) - \frac{dD_\Lambda^M}{d\Lambda} \quad (38)$$

$$\begin{aligned}
 & + 2(\dot{\Sigma}_\Lambda/S^2\mathcal{C} + S_\Lambda^{-1}\dot{S}_\Lambda)D_\Lambda^M - \dot{\Sigma}_\Lambda/S^2 (D_\Lambda^M)^2 \\
 & = -\frac{dD_\Lambda^M}{d\Lambda} + 2S_{2\Lambda}^{-1}\dot{S}_{2\Lambda}D_\Lambda^M - \dot{\Sigma}_\Lambda S^{-2} (D_\Lambda^M)^2
 \end{aligned}$$

where in the last line we have assumed $S = S_1 S_2$ and have chosen $S_1 = C_0^{-1} - \Sigma_\Lambda$. This choice of S_1 implies that the remaining factor S_2 represents the reamputation of the vertices, amputated by $C_\Lambda(C_0^{-1} - \Sigma_\Lambda)$. With this choice we also have

$$S^{-2}\dot{\Sigma}_\Lambda(\mathcal{C} - D_\Lambda^M) + \dot{S}_\Lambda S_\Lambda^{-1} = -\dot{\Sigma}_\Lambda S^{-2}D_\Lambda^M + \dot{S}_{2\Lambda}/S_{2\Lambda}$$

and

$$S^{-2}\dot{\Sigma}_\Lambda = S_{2\Lambda}^{-2}(\partial_\Lambda \Sigma_\Lambda)/(C_0^{-1} - \Sigma_\Lambda)^2 = S_{2\Lambda}^{-2}\partial_\Lambda(1/(C_0^{-1} - \Sigma_\Lambda)) = S_{2\Lambda}^{-2}\partial_\Lambda G_\Lambda$$

Introducing $\mathcal{Q}_\Lambda = S_{2\Lambda}^{-2}\partial_\Lambda G_\Lambda$, we obtain

$$\begin{aligned}
 \mathcal{Q}_\Lambda &= -\left[\frac{d}{d\Lambda}(\widetilde{D}_\Lambda - D_\Lambda^M) + 2D_\Lambda(D_\Lambda^M\mathcal{Q}_\Lambda - \dot{S}_{2\Lambda}S_{2\Lambda}^{-1})\right. \\
 &\quad \left.-(D_\Lambda^M)^2\mathcal{Q}_\Lambda + 2S_{2\Lambda}^{-1}\dot{S}_{2\Lambda}D_\Lambda^M\right]\widetilde{H}_4 \\
 &\quad - \sum_{m_1, m_2 \geq 4} \widetilde{H}_{m_1} \circ \left(\frac{dD_\Lambda^M}{d\Lambda} + \mathcal{Q}_\Lambda(D_\Lambda^M)^2 - 2S_{2\Lambda}^{-1}\dot{S}_{2\Lambda}D_\Lambda^M\right) \circ D_\Lambda^{\frac{m_1+m_2}{2}-2} \circ \widetilde{H}_{m_2} \\
 \dot{\mathcal{H}}_m &= \left[\frac{d}{d\Lambda}(\widetilde{D}_\Lambda - D_\Lambda^M) + 2D_\Lambda(\mathcal{Q}_\Lambda D_\Lambda^M - \dot{S}_{2\Lambda}S_{2\Lambda}^{-1})\right. \\
 &\quad \left.-(D_\Lambda^M)^2\mathcal{Q}_\Lambda + 2S_{2\Lambda}^{-1}\dot{S}_{2\Lambda}D_\Lambda^M\right]\widetilde{H}_{m+2} \\
 &\quad + \sum_{m_1, m_2 \geq 4} \widetilde{H}_{m_1} \circ \left(\frac{dD_\Lambda^M}{d\Lambda} + \mathcal{Q}_\Lambda(D_\Lambda^M)^2 - 2S_{2\Lambda}^{-1}\dot{S}_{2\Lambda}D_\Lambda^M\right) \circ D_\Lambda^{\frac{m_1+m_2-m}{2}-1} \circ \widetilde{H}_{m_2} \\
 &\quad + m\widetilde{H}_m(D_\Lambda^M\mathcal{Q}_\Lambda - \dot{S}_{2\Lambda}/S_{2\Lambda})
 \end{aligned} \tag{39}$$

For $D_\Lambda^M\mathcal{Q}_\Lambda - \dot{S}_{2\Lambda}/S_{2\Lambda} = 0$ we obtain equations

$$\begin{aligned}
 \mathcal{Q}_\Lambda &= -\left[\frac{d}{d\Lambda}(\widetilde{D}_\Lambda - D_\Lambda^M) + (D_\Lambda^M)^2\mathcal{Q}_\Lambda\right]\widetilde{H}_4 \\
 &\quad - \sum_{m_1, m_2 \geq 4} \widetilde{H}_{m_1} \circ \left(\frac{dD_\Lambda^M}{d\Lambda} - \mathcal{Q}_\Lambda(D_\Lambda^M)^2\right) \circ D_\Lambda^{\frac{m_1+m_2}{2}-2} \circ \widetilde{H}_{m_2} \\
 \dot{\mathcal{H}}_m &= \left[\frac{d}{d\Lambda}(\widetilde{D}_\Lambda - D_\Lambda^M) + (D_\Lambda^M)^2\mathcal{Q}_\Lambda\right]\widetilde{H}_{m+2} \\
 &\quad + \sum_{m_1, m_2 \geq 4} \widetilde{H}_{m_1} \circ \left(\frac{dD_\Lambda^M}{d\Lambda} - \mathcal{Q}_\Lambda(D_\Lambda^M)^2\right) \circ D_\Lambda^{\frac{m_1+m_2-m}{2}-1} \circ \widetilde{H}_{m_2}
 \end{aligned} \tag{40}$$

which coincide with the equations of Ref. [15]. Choosing the propagators as in Ref. [15], $D_M^\Lambda = \chi_{<,\Lambda}/\mathcal{A}_\Lambda$, $S_{2\Lambda} = 1/\mathcal{A}_\Lambda$, $\frac{d\mathcal{A}_\Lambda}{d\Lambda} = -\chi_{<,\Lambda}\mathcal{Q}_\Lambda$, we obtain

$$\begin{aligned}
 \mathcal{Q}_\Lambda &= -\sum_{m_1, m_2 \geq 4} \widetilde{H}_{m_1} \circ \frac{d\widetilde{D}_\Lambda}{d\Lambda} \circ \widetilde{D}_\Lambda^{\frac{m_1+m_2}{2}-2} \circ \widetilde{H}_{m_2}, \\
 \dot{\mathcal{H}}_m &= \sum_{m_1, m_2 \geq 4} \widetilde{H}_{m_1} \circ \frac{d\widetilde{D}_\Lambda}{d\Lambda} \circ \widetilde{D}_\Lambda^{\frac{m_1+m_2-m}{2}-1} \circ \widetilde{H}_{m_2},
 \end{aligned} \tag{41}$$

where the single-scale and Wick propagators are given by

$$\begin{aligned}\widetilde{D}_\Lambda &= \int_0^\Lambda d\Lambda' F_{\Lambda'}, \quad F_\Lambda = \frac{\dot{\chi}_{<,\Lambda}}{\mathcal{A}_\Lambda} \\ \mathcal{A}_\Lambda &= \frac{1}{C_0^{-1} - \Sigma_{\Lambda_0} + \int_{\Lambda_0}^{\Lambda} d\Lambda'' \chi_{<,\Lambda''} \mathcal{Q}_{\Lambda''}}.\end{aligned}\quad (42)$$

The relations (42) are not easily simplified further for the sharp cutoff because of the involved relation between $\mathcal{Q}_\Lambda$ and Σ_Λ , $\mathcal{Q}_\Lambda = [\mathcal{A}_\Lambda / (C_0^{-1} - \Sigma_\Lambda)]^2 \dot{\Sigma}_\Lambda$.

In the local scheme $\widetilde{D}_\Lambda = D_\Lambda^M$, also considered in Ref. [15], we have

$$\begin{aligned}\mathcal{Q}_\Lambda &= - \left(\frac{\chi_{<,\Lambda}}{\mathcal{A}_\Lambda} \right)^2 \mathcal{Q}_\Lambda \circ \widetilde{H}_4 \\ &\quad - \sum_{m_1, m_2 \geq 4} \widetilde{H}_{m_1} \circ \left(\frac{\dot{\chi}_{<,\Lambda}}{\mathcal{A}_\Lambda} \right) \circ \left(\frac{\chi_{<,\Lambda}}{\mathcal{A}_\Lambda} \right)^{\frac{m_1+m_2}{2}-2} \circ \widetilde{H}_{m_2} \\ \dot{\widetilde{H}}_m &= \left(\frac{\chi_{<,\Lambda}}{\mathcal{A}_\Lambda} \right)^2 \mathcal{Q}_\Lambda \circ \widetilde{H}_{m+2} \\ &\quad + \sum_{m_1, m_2 \geq 4} \widetilde{H}_{m_1} \circ \left(\frac{\dot{\chi}_{<,\Lambda}}{\mathcal{A}_\Lambda} \right) \circ \left(\frac{\chi_{<,\Lambda}}{\mathcal{A}_\Lambda} \right)^{\frac{m_1+m_2-m}{2}-1} \circ \widetilde{H}_{m_2}\end{aligned}\quad (43)$$

As well as the scheme (27), the latter equations may suffer for general many-body problems from divergencies of the convolution of $\widetilde{H}_{m \geq 4}$ with the square of propagator $\chi_{<,\Lambda}/\mathcal{A}_\Lambda$ and $\mathcal{Q}_\Lambda$, which are not cut in momentum space.

- [1] J. Polchinski, Nucl. Phys. B **231**, 269 (1984).
- [2] M. Salmhofer, *Comm. Math. Phys.* **194**, 249 (1998); *Renormalization* (Springer, Heidelberg, 1999).
- [3] N. Tetradis, J. Berges, C. Wetterich, Physics Reports, **363**, 223 (2002).
- [4] M. Salmhofer and C. Honerkamp, *Progr. Theor. Phys.* **105**, 1 (2001).
- [5] C. Honerkamp and M. Salmhofer, Phys. Rev. B **64**, 184516 (2001).
- [6] W. Metzner, M. Salmhofer, C. Honerkamp, V. Meden, and K. Schoenhammer, arXiv:1105.5289 (unpublished).
- [7] A. A. Katanin and A. P. Kampf, Phys. Rev. B **68**, 195101 (2003).
- [8] C. Honerkamp, D. Rohe, S. Andergassen, and T. Enss, Phys. Rev. B **70**, 235115 (2004).
- [9] P. A. Igoshev, V. Yu. Irkhin, and A. A. Katanin, Phys. Rev. B **83**, 245118 (2011).
- [10] A. Katanin, *Phys. Rev. B* **70**, 115109 (2004).
- [11] R. Hedden, V. Meden, Th. Pruschke, and K. Schoenhammer, *J. Phys.: Cond. Matt.* **16**, 5279 (2004); C. Karrasch, R. Hedden, R. Peters, Th. Pruschke, K. Schoenhammer, and V. Meden, *J. Phys.: Cond. Matt.* **20**, 345205 (2008).
- [12] M. Salmhofer, C. Honerkamp, W. Metzner, and O. Lauscher, *Progr. Theor. Phys.* **112**, 943 (2004); R. Gersch, C. Honerkamp, D. Rohe, and W. Metzner, *Eur. Phys. J. B.* **48**, 349 (2005); R. Gersch, C. Honerkamp, and W. Metzner, *New J. Phys.* **10**, 045003 (2008).
- [13] J. Reuther and P. Woelfle, *Phys. Rev. B* **81**, 144410 (2010); *Phys. Rev. B* **83**, 024402 (2011).
- [14] A. Katanin, *Phys. Rev. B* **79**, 235119 (2009).
- [15] M. Salmhofer, *Ann. der Phys.* **16**, 171 (2007).
- [16] R. Gersch, J. Reiss, and C. Honerkamp, *New J. Phys.* **8** 320 (2006).
- [17] V. Meden, the talk at the "Winter School on Renormalization Group Methods" in Brasilia, 2004, <http://web.physik.rwth-aachen.de/~meden/funRG/talk.pdf> (unpublished)